



Documentation Tips – Urology

Principal Diagnoses:

- A principal diagnosis (PDx) is the condition present on admission, requiring admission and treatment (may be more than one)
- Explain etiology where possible (*TURP performed due to recurrent prostatic CA*)

Secondary Diagnoses:

- Include *all* diagnoses present on admission which are in any way treated and/or monitored (re-exam, testing, meds, etc.)
 - This may be the only way to increase severity of illness
- Explain why the patient is at higher risk for surgery
- Utilize medical consults where appropriate to improve specificity of diagnoses, but document all diagnoses on *your* note as well

Conditions Arising During Hospitalization:

- Clearly describe your impression as to whether the condition is expected/anticipated (e.g. due to an underlying illness and/or commonly occurring and therefore *not* a complication) or a complication of care (e.g. unanticipated and/or outside of the expected post-op clinical course)
 - *“Acute Blood Loss Anemia, expected, secondary to bladder CA and cystectomy”*
 - *Acute blood loss anemia* is a CC
- Consider documenting a diagnosis any time you do something procedurally to a patient (suprapubic nephrostomy – *acute urinary retention*; CVC – hypovolemic shock)
- Beware of describing conditions in the post-op period as “post-op;” *post-op hypertension*, for example, codes as a complication – explain the etiology (e.g. *exacerbation of essential hypertension*)



Pearls for Urology Documentation

- Describe Your “Clinical Impression” of diagnosis/etiology
 - You can use words like probable, likely, suspect, etc.
- *Peritonitis* is an MCC
 - *Sepsis* (SIRS due to an infection) is an MCC, *or* may change the PDX if present on admission [*Urosepsis* = UTI to a coder]
- *Hydronephrosis* is a CC
- Acute Renal Failure/Acute Kidney Injury (AKI) – a CC
 - AKIN criteria - \uparrow in Cr by 0.3-0.5 above normal baseline = *St 1 AKI*, also a CC
 - Acute Renal *Insufficiency*, pre-renal azotemia, dehydration, etc = *low severity*
 - Acute Renal Failure with ATN (acute tubular necrosis) remains an MCC
- Document *severe malnutrition* – it not only adds severity as an MCC, it will likely prolong the post-op course thereby aligning the illness severity with length of stay
- Congestive Heart Failure
 - “CHF” no longer adds to severity
 - *Chronic systolic, diastolic or combined* heart failure adds severity via a minor comorbidity (CC)
 - *Acute systolic, diastolic or combined* heart failure adds severity via a *major* comorbidity (MCC)
- Describe *all* components of a surgical procedure as they often add complexity - which may also benefit your professional billing (“TURP *and* sphincterotomy of bladder neck due to BPH and bladder outlet obstruction due to recurrent infection”)
- Xray Reports
 - If an x-ray report demonstrates a clinical diagnosis, coders can’t code from an “interpreting physician’s report”
 - Document the condition in the progress notes
 - *e.g., Severe Interstitial Pulmonary Fibrosis*